

When to Consider SANCUSO[®] (Granisetron Transdermal System)

Due to its route of administration, SANCUSO may be an alternate option for patients who are unable to take or retain daily oral antiemetics^{1,2}

CANCER TYPE	TREATMENT REGIMENS						POSSIBLE COMPLICATIONS [†]	
	 Chemoradiation*	 Chemotherapy					 Difficulty Swallowing (e.g. oral mucositis) ^{12,13}	 Limited Gut Motility/ Absorption ^{14,15}
		Platinum-based (cisplatin, carboplatin)	Microtubule inhibitor (paclitaxel)	Antimetabolite (5-FU)	Anthracycline (doxorubicin, epirubicin)	Alkylating agent (cyclophos- phamide)		
 Head and neck ³	■	■	■	■			■	
Breast ⁴		■			■	■	■	
Lung ⁵	■	■					■	
Gastrointestinal ⁶	■	■		■	■		■	■
Gynecological ^{7,8}	■	■					■	■
Testicular ⁹		■					■	
Colorectal ^{10,11}	■	■		■			■	

*SANCUSO is not indicated for radiation-induced nausea and vomiting.

[†]Disease- or treatment-related complications that may make oral antiemetics difficult.

Indication

SANCUSO (Granisetron Transdermal System) is indicated for the prevention of nausea and vomiting in patients receiving moderately and/or highly emetogenic chemotherapy regimens of up to 5 consecutive days' duration.

Please see additional Important Safety Information on back and full Prescribing Information in pocket.

Think SANCUSO[®] (Granisetron Transdermal System)

If you check any of the boxes below, consider SANCUSO for your patient. Because it's a patch, SANCUSO may be appropriate for mechanically compromised* patients for whom daily oral antiemetics may not be an option.



Difficulty Swallowing

Possible causes:

- Side effects from chemotherapy with/without radiation, such as:
 - mouth sores¹³
 - oral mucositis^{12,13}
 - oral infections¹⁸
 - xerostomia/mouth dryness¹⁹
- Tumor-related dysmotility²⁰
- Chemoradiation^{21†}



Limited Gut Motility/Absorption

Possible causes:

- Side effects from chemotherapy with/without radiation, such as:
 - gastrointestinal mucositis²²
 - diarrhea²³
 - radiation-induced small bowel disease (radiation enteritis)^{24†}
 - bowel obstructions or perforations²⁵
- Chemoradiation^{26†}
- Opioids taken for pain management²⁷

* Mechanically compromised cancer patients include those with oral and/or gastrointestinal impairment associated with chemotherapy, with or without radiation,^{21,26,28,29} or tumor burden,^{27,30} who may be unable to take or retain oral antiemetics.²

† Sancuso is not indicated for radiation-induced nausea and vomiting.

Important Safety Information

SANCUSO is contraindicated in patients with known hypersensitivity to granisetron or to any of the components of the patch. Granisetron may mask a progressive ileus and/or gastric distention caused by the underlying condition. Mild application-site reactions have occurred; remove the patch if severe reaction or a generalized skin reaction occurs. Patients should avoid direct exposure of application site to natural or artificial sunlight by covering with clothing while wearing the patch and for 10 days after removing it. The most common adverse reaction in patients receiving SANCUSO is constipation (5.4%). SANCUSO contains granisetron. Healthcare professionals should avoid prescribing any additional products that contain granisetron. No clinically relevant drug interactions have been reported in clinical studies with SANCUSO. The development of serotonin syndrome has been reported with 5-HT₃ receptor antagonists. If symptoms of serotonin syndrome occur, discontinue SANCUSO and initiate supportive treatment. Patients should be informed of the increased risk of serotonin syndrome, especially if SANCUSO is used concomitantly with other serotonergic drugs.

References: 1. Sancuso [package insert]. Bridgewater, NJ; ProStrakan Inc 2014. 2. Schulmeister L. Granisetron transdermal system: a new option to help prevent chemotherapy-induced nausea and vomiting. *Clin J Oncol Nurs.* 2009;13(6):711-714. 3. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Head and neck cancers. V.2.2014. 4. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Breast cancer. V.3.2014. 5. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Non-small cell lung cancer. V.1.2015. 6. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Gastric cancer. V.1.2014. 7. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Cervical cancer. V.1.2015. 8. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Ovarian cancer: including fallopian tube cancer and primary peritoneal cancer. V.3.2014. 9. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Testicular cancer. V.1.2014. 10. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Colon cancer. V.1.2015. 11. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Rectal cancer. V.1.2015. 12. Lalla RV, Bowen J, Barasch A, et al. MASCC/ISO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer.* 2014;120:1453-1461. 13. Cawley MM, Benson LM. Current trends in managing oral mucositis. *Clin J Oncol Nurs.* 2005;9(5):584-592. 14. Di Lorenzo C, Youssef NN. Diagnosis and management of intestinal motility disorders. *Semin Pediatr Surg.* 2010;19:50-58. 15. Keller J, Layer P. Intestinal and anorectal motility and functional disorders. *Best Pract Res Clin Gastroenterol.* 2009;23:407-423. 16. Tuca A. Use of granisetron transdermal system in the prevention of chemotherapy-induced nausea and vomiting: a review. *Cancer Manag Res.* 2009;2:1-12. 17. Boccia RV. Chemotherapy-induced nausea and vomiting: identifying and addressing unmet needs. *JCOM.* 2013;20(8):377-384. 18. Lalla RV, Latorue MC, Hong CH, et al. A systematic review of oral fungal infections in patients receiving cancer therapy. *Support Care Cancer.* 2010;18(8):985-992. 19. Jensen SB, Pedersen AML, Vissink A, et al. A systematic review of salivary gland hypofunction and xerostomia induced by cancer therapies: prevalence, severity and impact on quality of life. *Support Care Cancer.* 2010;18:1039-1060. 20. National Comprehensive Cancer Network website. NCCN Clinical Practice Guidelines in Oncology: Esophageal and esophagogastric junction cancers. V.3.2015. 21. National Cancer Institute website. Oral complications of chemotherapy and head/neck radiation. <http://www.cancer.gov/cancertopics/pdq/supportivecare/oralcomplications/HealthProfessional>. Accessed on August 20, 2014. 22. Keefe DM. Intestinal mucositis: mechanisms and management. *Curr Opin Oncol.* 2007;19:323-327. 23. Muehlbauer PM, Thorpe D, Davis A, et al. Putting evidence into practice: evidence-based interventions to prevent, manage, an treat chemotherapy- and radiotherapy-induced diarrhea. *Clin J Oncol Nurs.* 2009;13(3):336-341. 24. Stacey R, Green JT. Radiation-induced small bowel disease: latest developments and clinical guidance. *Ther Adv Chronic Dis.* 2014;5(1):15-29. 25. Ripamonti CI, Easson AM, Gerdes H. Management of malignant bowel obstruction. *Eur J Cancer.* 2008;44:1105-1115. 26. National Cancer Institute website. Nausea and Vomiting (PDQ[®]) Radiation Therapy. <http://www.cancer.gov/cancertopics/pdq/supportivecare/nausea/HealthProfessional/page9>. Accessed on September 16, 2014. 27. National Cancer Institute website. Nausea and Vomiting (PDQ[®]) Nausea, Vomiting (Emesis), Constipation, and Bowel Obstruction in Advanced Cancer. <http://www.cancer.gov/cancertopics/pdq/supportivecare/nausea/HealthProfessional/page7>. Accessed on September 16, 2014. 28. Raber-Durlacher JE, Brennan MT, Verdonck-de Leeuw IM, et al. Swallowing dysfunction in cancer patients. *Support Care Cancer.* 2012;20:433-443. 29. Roeder F, Nicolay NH, Nguyen T, et al. Intensity modulated radiotherapy (IMRT) with concurrent chemotherapy as definitive treatment of locally advanced esophageal cancer. *Radiat Oncol.* 2014;9:191. 30. Hursti TJ, Avall-Lundqvist E, Borjeson S, et al. Impact of tumor burden on chemotherapy-induced nausea and vomiting. *Br J Cancer.* 1996;74(7):1114-1119.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Sancuso (Granisetron Transdermal System) safely and effectively. See full prescribing information for Sancuso.

Sancuso (Granisetron Transdermal System) Initial U.S. Approval: 2008

-----RECENT MAJOR CHANGES-----

Warnings and Precautions, Serotonin Syndrome (5.4) 09/2014

-----INDICATIONS AND USAGE-----

Sancuso is a serotonin subtype 3 (5-HT₃) receptor antagonist indicated for the prevention of nausea and vomiting in patients receiving moderately and/or highly emetogenic chemotherapy for up to 5 consecutive days. (1)

----DOSAGE AND ADMINISTRATION----

Apply a single transdermal system (patch) to the upper outer arm a minimum of 24 hours before chemotherapy. The patch may be applied up to a maximum of 48 hours before chemotherapy as appropriate. Remove the patch a minimum of 24 hours after completion of chemotherapy. The patch can be worn for up to 7 days depending on the duration of the chemotherapy regimen. (2)

---DOSAGE FORMS AND STRENGTHS---

52 cm² patch containing 34.3 mg of granisetron delivering 3.1 mg per 24 hours. (3)

-----CONTRAINDICATIONS-----

Known hypersensitivity to granisetron or to any of the components of the patch. (4)

----WARNINGS AND PRECAUTIONS----

- Granisetron may mask a progressive ileus and/or gastric distention caused by the underlying condition. (5.1)
- Mild application site reactions have occurred; remove patch if severe reactions or a generalized skin reaction occur. (5.2)
- Avoid direct exposure of application site to natural or artificial sunlight by covering with clothing while wearing the patch and for 10 days after removing it. (5.3)
- Serotonin syndrome has been reported with 5-HT₃ receptor antagonists alone but particularly with concomitant use of serotonergic drugs. (5.4)

-----ADVERSE REACTIONS-----

The most common adverse reaction is constipation. (6)

To report SUSPECTED ADVERSE REACTIONS, contact ProStrakan Inc. at 1-800-SANCUSO and www.sancuso.com, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

- No clinically relevant drug interactions have been reported in clinical studies with Sancuso. (7)
- Use during pregnancy only if clearly needed. (8.1)
- Use caution when administering to nursing women. (8.3)

- Safety and effectiveness in pediatric patients have not been established. (8.4)
- Clinical studies of Sancuso did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger patients. (8.5)

----USE IN SPECIFIC POPULATIONS----

Use in pregnancy only if clearly needed. (8.1)

Use caution when administering to nursing women. (8.3)

Safety and effectiveness in pediatric patients have not been established. (8.4)

Clinical studies of Sancuso did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger patients. (8.5)

See **17 for PATIENT COUNSELING INFORMATION** and **FDA-approved patient labeling.**

Revised: 09/2014

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Sancuso® (Granisetron Transdermal System) is indicated for the prevention of nausea and vomiting in patients receiving moderately and/or highly emetogenic chemotherapy regimens of up to 5 consecutive days duration.

2 DOSAGE AND ADMINISTRATION

The transdermal system (patch) should be applied to clean, dry, intact healthy skin on the upper outer arm. Sancuso should not be placed on skin that is red, irritated or damaged.

Each patch is packed in a pouch and should be applied directly after the pouch has been opened. The patch should not be cut into pieces.

2.1 Adults

Apply a single patch to the upper outer arm a minimum of 24 hours before chemotherapy. The patch may be applied up to a maximum of 48 hours before chemotherapy as appropriate. Remove the patch a minimum of 24 hours after completion of chemotherapy. The patch can be worn for up to 7 days depending on the duration of the chemotherapy regimen.

3 DOSAGE FORMS AND STRENGTHS

Sancuso is a 52 cm² patch containing 34.3 mg of granisetron. The patch releases 3.1 mg of granisetron per 24 hours for up to 7 days.

4 CONTRAINDICATIONS

Sancuso is contraindicated in patients with known hypersensitivity to granisetron or to any of the components of the patch.

5 WARNINGS AND PRECAUTIONS

5.1 Gastrointestinal
The use of granisetron in patients may mask a progressive ileus and/or gastric distention caused by the underlying condition.

5.2 Skin Reactions

In clinical trials with Sancuso, application site reactions were reported which were generally mild in intensity and did not lead to discontinuation of use. The incidence of reactions was comparable with placebo.

If severe reactions, or a generalized skin reaction occur (e.g. allergic rash, including erythematous, macular, papular rash or pruritus), the patch must be removed.

5.3 Exposure to Sunlight

Granisetron may be affected by direct natural or artificial sunlight. Patients must be advised to cover the patch application site, e.g. with clothing, if there is a risk of exposure to sunlight throughout the period of wear and for 10 days following its removal because of a potential skin reaction (see Section 13.2).

5.4 Serotonin Syndrome

The development of serotonin syndrome has been reported with 5-HT₃ receptor antagonists. Most reports have been associated with concomitant use of serotonergic drugs

(e.g., selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), monoamine oxidase inhibitors, mirtazapine, fentanyl, lithium, tramadol, and intravenous methylene blue). Some of the reported cases were fatal. Serotonin syndrome occurring with overdose of another 5-HT₃ receptor antagonist alone has also been reported.

The majority of reports of serotonin syndrome related to 5-HT₃ receptor antagonist use occurred in a post-anesthesia care unit or an infusion center.

Symptoms associated with serotonin syndrome may include the following combination of signs and symptoms: mental status changes

(e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, with or without gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Patients should be monitored for the emergence of serotonin syndrome, especially with concomitant use of Sancuso and other serotonergic drugs. If symptoms of serotonin syndrome occur, discontinue Sancuso and initiate supportive treatment. Patients should be informed of the increased risk of serotonin syndrome, especially if Sancuso is used concomitantly with other serotonergic drugs. [see *Drug Interactions (7), Patient Counseling Information (17.4).*]

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

The safety of Sancuso was evaluated in a total of 404 patients undergoing chemotherapy who participated in two double-blind, comparator studies with patch treatment durations of up to 7 days. The control groups included a total of 406 patients who received a daily dose of 2 mg oral granisetron, for 1 to 5 days.

Adverse reactions considered by the investigators as drug-related occurred in 8.7% (35/404) of patients receiving Sancuso and 7.1% (29/406) of patients receiving oral granisetron. The most common adverse reaction was constipation that occurred in 5.4% of patients in the Sancuso group and 3.0% of patients in the oral granisetron group. Table 1 lists the treatment emergent adverse reactions that occurred in at least 3% of patients treated with Sancuso or oral granisetron.

Table 1: Incidence of Adverse Reactions in Double-Blind, Active Comparator Controlled Studies in Cancer Patients Receiving Chemotherapy (Events ≥ 3% in either group)

Body System Preferred Term	Sancuso TDS N=404 (%)	Oral granisetron N=406 (%)
Gastrointestinal disorders		
Constipation	5.4	3.0
Nervous system disorders		
Headache	0.7	3.0

5-HT₃ receptor antagonists, such as granisetron, may be associated with arrhythmias or ECG abnormalities. Three ECGs were performed on 588 randomized patients in the Phase 3 study: at baseline before treatment, the first day of chemotherapy, and 5 to 7 days after starting chemotherapy. QTcF prolongation greater than 450 milliseconds was seen in a total of 11 (1.9%) patients after receiving granisetron, 8 (2.7%) on oral granisetron and 3 (1.1%) on the patch. No new QTcF prolongation greater than 480 milliseconds was observed in any patient in this study. No arrhythmias were detected in this study.

6.2 Granisetron Experience
Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse events reported in clinical trials with other formulations of granisetron include the following: **Gastrointestinal:** abdominal pain, diarrhea, constipation, elevation of ALT and AST levels, nausea and vomiting

Cardiovascular: hypertension, hypotension, angina pectoris, atrial fibrillation and syncope have been observed rarely

Central Nervous System: dizziness, insomnia, headache, anxiety, somnolence and asthenia

Hypersensitivity: rare cases of hypersensitivity reactions, sometimes severe (e.g. anaphylaxis, shortness of breath, hypotension, urticaria) have been reported

Other: fever; events often associated with chemotherapy have also been reported: leucopenia, decreased appetite, anemia, alopecia, thrombocytopenia.

7 DRUG INTERACTIONS

Granisetron does not induce or inhibit the cytochrome P-450 drug-metabolizing enzyme system *in vitro*. There have been no definitive drug-drug interaction studies to examine pharmacokinetic or pharmacodynamic interaction with other drugs. However, in humans, granisetron hydrochloride injection has been safely administered with drugs representing benzodiazepines, neuroleptics and anti-ulcer medications commonly prescribed with antiemetic treatments. Granisetron hydrochloride injection also does not appear to interact with emetogenic cancer therapies. In agreement with these data, no clinically relevant drug interactions have been reported in clinical studies with Sancuso.

Because granisetron is metabolized by hepatic cytochrome P-450 drug-metabolizing enzymes (CYP1A1 and CYP3A4), inducers or inhibitors of these enzymes may change the clearance and hence, the half-life of granisetron. In addition, the activity of the cytochrome P-450 subfamily 3A4 (involved in the metabolism of some of the main narcotic analgesic agents) is not modified by granisetron hydrochloride *in vitro*. In *in vitro* human microsomal studies, ketoconazole inhibited ring oxidation of granisetron hydrochloride. However, the clinical significance of *in vivo* pharmacokinetic interactions with ketoconazole is not known. In a human pharmacokinetic study, hepatic enzyme induction with phenobarbital resulted in a 25% increase in total plasma clearance of intravenous granisetron hydrochloride. The clinical significance of this change is not known.

Serotonin syndrome (including altered mental status, autonomic instability, and neuromuscular symptoms) has been described following the concomitant use of 5-HT₃ receptor antagonists and other serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs) and serotonin and norepinephrine reuptake inhibitors (SNRIs) [see Warnings and Precautions (5.4)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B
Reproduction studies with granisetron hydrochloride have been performed in pregnant rats at intravenous doses up to 9 mg/kg/day (54 mg/m²/day, about 24 times the recommended human dose delivered by the Sancuso patch, based on body surface area) and oral doses up to 125 mg/kg/day (750 mg/m²/day, about 326 times the recommended human dose with Sancuso based on body surface area). Reproduction studies have been performed in pregnant rabbits at intravenous doses up to 3 mg/kg/day (36 mg/m²/day, about 16 times the human dose with Sancuso based on body surface area) and at oral doses up to

32 mg/kg/day (384 mg/m²/day, about 167 times the human dose with Sancuso based on body surface area). These studies did not reveal any evidence of impaired fertility or harm to the fetus due to granisetron. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, Sancuso should be used during pregnancy only if clearly needed.

8.3 Nursing Mothers

It is not known whether granisetron is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Sancuso is administered to a nursing woman.

8.4 Pediatric Use

Safety and effectiveness of Sancuso in pediatric patients under 18 years of age have not been established.

8.5 Geriatric Use

Clinical studies of Sancuso did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, cautious treatment selection for an elderly patient is prudent because of the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Renal Failure or Hepatically-Impaired Patients

Although no studies have been performed to investigate the pharmacokinetics of Sancuso in patients with renal or hepatic impairment, pharmacokinetic information is available for intravenous granisetron (see CLINICAL PHARMACOLOGY: Pharmacokinetics 12.3).

10 OVERDOSAGE

There is no specific antidote for granisetron overdose. In the case of overdose, symptomatic treatment should be given. Overdose of up to 38.5 mg of granisetron hydrochloride, as a single intravenous injection, has been reported without symptoms or only the occurrence of a slight headache. In clinical trials there were no reported cases of overdose with Sancuso.

11 DESCRIPTION

Sancuso contains granisetron, which is an anti-nauseant and antiemetic agent. Chemically it is 1-methyl-N-[(1R,3r,5S)-9-methyl-9-azabicyclo[3.3.1]non-3-yl]-1H-indazole-3-carboxamide with a molecular weight of 312.4. Its empirical formula is C₁₈H₂₄N₄O, while its chemical structure is:

Granisetron crosses intact skin into the systemic circulation by a passive diffusion process.

Following a 7-day application of Sancuso in 24 healthy subjects, high inter-subject variability in systemic exposure was observed. Maximal concentration was reached at approximately 48 hours (range: 24-168 hours) following patch application. Mean C_{max} was 5.0 ng/mL (CV: 170%) and mean AUC_{0-168h} was 527 ng-hr/mL (CV:173%).

Based on the measure of residual content of the patch after removal, approximately 66% (SD: ± 10.9) of granisetron is delivered following patch application for 7 days.

Granisetron is a white to off-white solid that is insoluble in water. Sancuso is a thin, translucent, matrix-type transdermal patch that is rectangular-shaped with rounded corners, consisting of a backing, the drug matrix and a release liner.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Granisetron is a selective 5-hydroxytryptamine₃ (5-HT₃) receptor antagonist with little or no affinity for other serotonin receptors, including 5-HT₁, 5-HT_{1A}, 5-HT_{1B/C}, 5-HT₂, for alpha₁-, alpha₂-, or beta-adrenoreceptors; for dopamine-D₂; or for histamine-H₁; benzodiazepine; pirototoxin or opioid receptors.

Serotonin receptors of the 5-HT₃ type are located peripherally on vagal nerve terminals and centrally in the chemoreceptor trigger zone of the area postrema. During chemotherapy that induces vomiting, mucosal enterochromaffin cells release serotonin, which stimulates 5-HT₃ receptors. This evokes vagal afferent discharge, inducing vomiting. Animal studies demonstrate that, in binding to 5-HT₃ receptors, granisetron blocks serotonin stimulation and subsequent vomiting after emetogenic stimuli such as cisplatin. In the ferret animal model, a single granisetron injection prevented vomiting due to high-dose cisplatin or arrested vomiting within 5 to 30 seconds.

12.2 Pharmacodynamics

The effect of granisetron on QTc prolongation was evaluated in a randomized, single-blind, positive (moxifloxacin 400 mg) - and placebo controlled parallel study in healthy subjects. A total of 240 subjects were administered Sancuso patch, intravenous granisetron (10 mcg/kg over 30 seconds). In a study with demonstrated ability to detect small effects, the upper bound of the 90% confidence interval for the largest placebo adjusted, baseline corrected QTc based on Fridericia correction method (QTcF) for Sancuso was below 10 ms, the threshold for regulatory concern.

No evidence of an effect on plasma prolactin or aldosterone concentrations has been found in studies using granisetron. The effect on oro-cecal transit time following application of Sancuso has not been studied. Granisetron hydrochloride injection exhibited no effect on oro-cecal transit time in healthy subjects given a single intravenous infusion of 50 mcg/kg or 200 mcg/kg. Single and multiple oral doses of granisetron hydrochloride slowed colonic transit in healthy subjects.

Elderly, and Renal or Hepatic Impairment
Although no studies have been performed to investigate the pharmacokinetics of Sancuso in elderly subjects, and in patients with renal or hepatic impairment, the following pharmacokinetic information is available for intravenous granisetron.

In the elderly, and in patients with renal failure or hepatic impairment, the pharmacokinetics of granisetron were determined following a single 40 mcg/kg intravenous dose of granisetron hydrochloride. In elderly volunteers (mean age 71 years) pharmacokinetic parameters following a single 40 mcg/kg intravenous dose of granisetron hydrochloride, lower clearance and longer half-life were observed compared to younger healthy volunteers.

Renal Failure Patients
Total clearance of granisetron was not affected in patients with severe renal failure who received a single 40 mcg/kg intravenous dose of granisetron hydrochloride.

Hepatically-Impaired Patients
In patients with hepatic impairment due to neoplastic liver involvement, total plasma clearance following a single 40 mcg/kg intravenous dose of granisetron hydrochloride was approximately halved compared to patients without hepatic impairment. Given the wide variability in pharmacokinetic parameters of granisetron and the good tolerance of doses well above the recommended dose, dose adjustment in patients with hepatic functional impairment is not necessary.

Absorption
Granisetron crosses intact skin into the systemic circulation by a passive diffusion process.

Following a 7-day application of Sancuso in 24 healthy subjects, high inter-subject variability in systemic exposure was observed. Maximal concentration was reached at approximately 48 hours (range: 24-168 hours) following patch application. Mean C_{max} was 5.0 ng/mL (CV: 170%) and mean AUC_{0-168h} was 527 ng-hr/mL (CV:173%).

Based on the measure of residual content of the patch after removal, approximately 66% (SD: ± 10.9) of granisetron is delivered following patch application for 7 days.

Plasma protein binding is approximately 65%. Granisetron distributes freely between plasma and red blood cells.

Elimination
Clearance is predominantly by hepatic metabolism. Based on a study with intravenous injection, approximately 12% of the dose is excreted unchanged in the urine of healthy subjects in 48 hours. The remainder of the dose is excreted as metabolites, 49% in the urine, and 34% in the feces.

Subpopulations
Gender
There is evidence to suggest that female subjects had higher granisetron concentrations than males following patch application. However, no statistically significant difference in clinical efficacy outcome was observed between genders.

Patient Information

Sancuso® [san-KOO-so] (Granisetron Transdermal System)

IMPORTANT: For skin use only

Read the Patient Information that comes with Sancuso before you start using it and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment. If you have any questions about Sancuso, ask your healthcare provider.

What is Sancuso?

Sancuso is a prescription medicine used to prevent nausea and vomiting in people receiving some types of chemotherapy treatment. Sancuso is a skin patch that slowly releases the medicine contained in the adhesive (glue), through clean and intact skin areas into your bloodstream while you wear the patch.

Important: Sancuso contains granisetron, the same medicine in Kytril. Do not take Kytril at the same time you use Sancuso unless your healthcare provider tells you it is alright.

How should I store Sancuso?

- Keep Sancuso in the package it comes in.
- Store Sancuso at 20-25°C (68-77°F).

Keep Sancuso out of the reach of children.

General information about Sancuso

Medicines are sometimes prescribed for conditions that are not mentioned in Patient Information leaflets. Do not use Sancuso for a condition for which it is not prescribed. Do not give Sancuso to other people, even if they have the same symptoms you have. It may harm them.

Who should not use Sancuso?

Do not use Sancuso if you are allergic to any of the ingredients in Sancuso. See the end of this leaflet for a list of ingredients in Sancuso.

What should I tell my healthcare provider before using Sancuso?

Tell your healthcare provider about all your medical conditions, including if you:

- are allergic to medical adhesive tape, adhesive dressings or other skin patches.
- have pain or swelling in your stomach area (abdomen).
- are pregnant. It is not known if Sancuso will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant.
- are breast-feeding or plan to breast-feed. It is not known if Sancuso passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and non-prescription medicines, vitamins and herbal supplements. Other medicines may affect how Sancuso works. Sancuso may also affect how other medicines work.

How should Sancuso be used?

Use Sancuso exactly as prescribed. See the detailed Patient Instructions for Applying Sancuso at the end of this Patient Information leaflet.

What should I avoid while using Sancuso?

Avoid sunlight. The medicine in Sancuso (granisetron) may not work as well and/or may affect your skin if exposed to direct sunlight or the light from sunlamps or tanning beds. It is important to do the following:

- While you wear the patch, keep it covered with clothing if you will be in sunlight or near a sunlamp, including tanning beds.
- Keep the skin where Sancuso was applied covered for another 10 days after the patch is taken off to protect from exposure to direct sunlight.

What are the possible side effects of Sancuso?

Sancuso can cause serious side effects:

- Using Sancuso may make it harder to identify certain stomach and bowel problems that are from other causes.** Tell your healthcare provider if you have any stomach area (abdominal) pain or swelling while using Sancuso.
- Skin reactions.** Skin reactions can happen just at the patch application site or outside the patch application site. Tell your healthcare provider if you get any redness, rashes, bumps, blisters or itching at the patch application site, and especially if they spread outside the place where the patch was or if they appear outside the patch application site. You may need to stop using Sancuso.

Common side effects of Sancuso are:

- constipation
- headache.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Sancuso. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Sancuso?

- Keep Sancuso in the package it comes in.
- Store Sancuso at 20-25°C (68-77°F).

Keep Sancuso out of the reach of children.

General information about Sancuso

Medicines are sometimes prescribed for conditions that are not mentioned in Patient Information leaflets. Do not use Sancuso for a condition for which it is not prescribed. Do not give Sancuso to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information leaflet summarizes the most important information about Sancuso. If you would like more information, talk to your healthcare provider. You can ask your pharmacist or healthcare provider for information about Sancuso that is written for health professionals.

For more information, go to www.sancuso.com or call 1-800-SANCUSO.

Patent Instructions for Applying Sancuso

When do I apply the Sancuso patch?

- Apply Sancuso at least 1 day (24 hours) before your scheduled chemotherapy treatment.
- You may apply Sancuso up to 2 days (48 hours) before your scheduled chemotherapy.
- Wear the patch all the time during your chemotherapy.
- Sancuso may be worn for up to 7 days, depending on how long your chemotherapy treatment lasts (up to 5 days).
- Remove the patch at least 1 day (24 hours) after your chemotherapy is finished.
- Keep the patch covered, such as under clothing, while you are wearing it to avoid a skin reaction to sunlight or sunlamps. Keep the skin where Sancuso was applied (application site) covered up for another 10 days after the patch is taken off to prevent a skin reaction. See "What should I avoid while using Sancuso?"

Where do I apply the Sancuso patch?

- Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.

Apply Sancuso to a clean, dry, healthy area of skin on the


How do I apply the Sancuso patch?

The Sancuso patch comes inside a pouch which is inside the carton.

- Do not remove the patch from the pouch until you are ready to use it.
- Do not cut the Sancuso patch into pieces.**
- Remove the pouch from the carton.




- Tear the pouch open using the slit provided, and remove the patch. Each pouch contains one Sancuso patch stuck onto a rigid plastic film, and a separate thin, clear protective liner.




- Remove the thin, clear protective liner to expose the printed side of the patch. Throw away the liner. The protective liner is only included in the pouch to separate the patch from the inside of the pouch, and is not part of the patch.



- The unprinted, sticky side of the patch is covered by a two-piece rigid plastic film. Bend the patch in the middle and remove one half of the rigid plastic film. Be careful not to stick the patch to itself and avoid touching the sticky side of the patch.



- While holding the remaining half of the rigid plastic film, apply the patch to your skin. Remove the second half of the rigid plastic film and press the whole patch firmly in place with your fingers and smooth down. Press firmly making sure it sticks well to the skin, especially around the edges.



- Wash your hands right away after applying the patch to remove any medicine that may have stuck to your fingers.
- Keep the patch in place for the whole time you are having chemotherapy. Remove the patch at least 1 day (24 hours) after your chemotherapy is finished. The patch can be worn for up to 7 days, depending on the number of days your chemotherapy treatment lasts.
- Do not re-use the patch after you remove it. See below for instructions on the right way to remove and throw away the patch.

What to do if the Sancuso patch does not stick well?

If the patch does not stick well, you may use surgical bandages or medical adhesive tape to keep the patch in place. Place tape or bandages on the edges of the patch. Do not completely cover the patch with bandages or tape and do not wrap completely around your arm. If the patch comes more than half off or it becomes damaged see your healthcare provider.

Can I bathe or shower while wearing Sancuso?

You can continue to shower and wash normally while wearing the Sancuso patch. It is not known how other activities, for example swimming, strenuous exercise or using a sauna or whirlpool, may affect Sancuso. Avoid these activities while wearing Sancuso.

How do I remove and dispose of Sancuso?

- When you remove the patch, peel it off gently.
- The used patch will still contain some of the medicine. After removing the used Sancuso patch, fold it in half so that the sticky side sticks to itself. Throw away the Sancuso patch in the garbage, out of the reach of children and pets. Do not re-use the patch.
- After removing the patch you may find some adhesive is left on your skin. Gently wash the area with soap and water to remove it. Do not use alcohol or other dissolving liquids, such as nail polish remover. These may cause skin irritation.
- Wash your hands after handling the patch.
- You may see mild redness on the skin where the patch is removed. This redness should go away within three days. If redness continues, tell your healthcare provider.

What are the ingredients in Sancuso?

Active ingredient: granisetron. Inactive ingredients: acrylate-vinylacetate copolymer, polyester, titanium dioxide, polyamide resin and polyethylene wax.

Manufactured by:

3M Drug Delivery Systems, St. Paul, MN 55107

Manufactured for:

ProStrakan Inc., Bridgewater, NJ 08807

Copyright © 2013 ProStrakan Inc. All rights reserved.

Sancuso and ProStrakan are trademarks owned by the ProStrakan group of companies.

Revised: January 2013

NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis and Impairment of Fertility

In a 24-month carcinogenicity study, rats were treated orally with granisetron 1, 5 or 50 mg/kg/day (6, 30 or 300 mg/m²/day). The 50 mg/kg/day dose was reduced to 25 mg/kg/day (150 mg/m²/day) during week 59 due to toxicity. For a 50 kg person of average height (1.46 m² body surface area), these doses represent about 2.6, 13 and 65 times the recommended clinical dose (3.1 mg/day, 2.3 mg/m²/day, delivered by the Sancuso patch, on a body surface area basis). There was a statistically significant increase in the incidence of hepatocellular carcinomas and adenomas in males treated with 5 mg/kg/day (30 mg/m²/day, about 13 times the recommended human dose with Sancuso, on a body surface area basis) and above, and in females treated with 25 mg/kg/day (150 mg/m²/day, about 65 times the recommended human dose with Sancuso, on a body surface area basis). No increase in liver tumors was observed at a dose of 1 mg/kg/day (6 mg/m²/day, about 2.6 times the recommended human dose with Sancuso, on a body surface area basis) in males and 5 mg/kg/day (30 mg/m²/day, about 13 times the recommended human dose with Sancuso, on a body surface area basis) in females.

In a 12-month oral toxicity study, treatment with granisetron 100 mg/kg/day (600 mg/m²/day), about 261 times the recommended human dose with Sancuso, on a body surface area basis) produced hepatocellular adenomas in male and female rats while no such tumors were found in the control rats. A 24-month mouse carcinogenicity study of granisetron did not show a statistically significant increase in tumor incidence, but the study was not conclusive.

Because of the tumor findings in rat studies, Sancuso should be prescribed only at the dose and for the indication recommended (see INDICATIONS AND USAGE, and DOSAGE AND ADMINISTRATION).

Granisetron was not mutagenic in an *in vitro* Ames test and mouse lymphoma cell forward mutation assay, and *in vivo* mouse micronucleus test and *in vitro* and *ex vivo* rat hepatocyte UDS assays. It, however, produced a significant increase in UDS in HeLa cells *in vitro* and a significant increased incidence of cells with polyploidy in an *in vitro* human lymphocyte chromosomal aberration test.

Granisetron at subcutaneous doses up to 6 mg/kg/day (36 mg/m²/day, about 16 times the recommended human dose of Sancuso, on a body surface area basis), and oral doses up to 100 mg/kg/day (600 mg/m²/day, about 261 times the recommended human dose of Sancuso, on a body surface area basis) was found to have no effect on fertility and reproductive performance of male and female rats.

Granisetron was not phototoxic when tested *in vitro* in a Chinese hamster ovary (CHO) cell line, at 200 and 300 mcg/ml, granisetron increased the percentage of cells with chromosomal aberration following photoirradiation.

Granisetron was not phototoxic when tested *in vitro* in a mouse fibroblast cell line. When tested *in vivo* in guinea-pigs, Sancuso patches did not show any potential for photirritation or photosensitivity. No phototoxicity studies have been performed in humans.

14. CLINICAL STUDIES

The effectiveness of Sancuso in the prevention of chemotherapy-induced nausea and vomiting (CINV) was evaluated in a Phase 3 randomized, parallel group, double-blind, double-dummy study conducted in the U.S. and abroad. The study compared the efficacy, tolerability and safety of Sancuso with that of 2 mg oral granisetron once daily in the prevention of nausea and vomiting in a total of 641 patients receiving multi-day chemotherapy.

The population randomized into the trial included 48% males and 52% females aged 16 to 86 years receiving moderately (ME) or highly emetogenic (HE) multi-day chemotherapy. Seventy-eight (78%) of patients were White, 12% Asian, 10% Hispanic/Latino and 0% Black.

The granisetron patch was applied 24 to 48 hours before the first dose of chemotherapy, and kept in place for 7 days. Oral granisetron was administered daily for the duration of the chemotherapy regimen, one hour before each dose of chemotherapy. Efficacy was assessed from the first administration until 24 hours after the start of the last day's administration of the chemotherapy regimen.

The primary endpoint of the trial was the proportion of patients achieving no vomiting and/or retching, no more than mild nausea and no rescue medication from the first administration until 24 hours after the start of the last day's administration of multi-day chemotherapy. Using this definition, the effect of Sancuso was established in 60.2% of patients in the Sancuso arm and 64.8% of patients receiving oral granisetron (difference -4.89%; 95% confidence interval -12.91% to +3.13%).

An assessment of patch adhesion in 621 patients receiving either active or placebo patches showed that less than 1% of patches became detached over the course of the 7 day period of patch application.

16. HOW SUPPLIED/STORAGE AND HANDLING

Sancuso (Granisetron Transdermal System) is supplied as a 52 cm² patch containing 34.3 mg of granisetron. Each patch is printed on one side with the words "Granisetron 3.1 mg/24 hours". Each patch is packaged in a separate sealed foil-lined plastic pouch.

Sancuso is available in packages of 1 (NDC 42747-726-01) patch.

Store at 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F). [see USP Controlled Room Temperature].

Sancuso should be stored in the original packaging.

17. PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (17.5)

17.1 Gastrointestinal

Because the use of granisetron may mask a progressive ileus and/or gastric distention caused by the underlying condition, patients should be instructed to tell their physician if they have pain or swelling in their abdomen.

17.2 Skin Reactions

Patients should be instructed to remove the patch if they have a severe skin reaction, or a generalized skin reaction (e.g. allergic rash, including erythematous, macular, papular rash or pruritus).

When patients remove the patch, they should be instructed to peel it off gently.

17.3 Exposure to Sunlight

Granisetron may be degraded by direct sunlight or exposure to sunlamps. In addition, an *in vitro* study using Chinese hamster ovary cells suggests that granisetron has the potential for photogenotoxicity (see Section 13.2).

Patients must be advised to cover the patch application site, e.g. with clothing, if there is a risk of exposure to sunlight or sunlamps throughout the period of wear and for 10 days following its removal.

17.4 Serotonin Syndrome

Advise patients of the possibility of serotonin syndrome with concomitant use of Sancuso and another serotonergic agent such as medications to treat depression and migraines. Advise patients to seek immediate medical attention if the following symptoms occur: changes in mental status, autonomic instability, neuromuscular symptoms, with or without gastrointestinal symptoms.

Rx only

Manufactured by:

3M Drug Delivery Systems, St. Paul, MN 55107

Manufactured for:

ProStrakan Inc.,
Bridgewater, NJ, 08807



Rev. 09/14

Copyright © 2014, ProStrakan Inc. All rights reserved.

Sancuso and ProStrakan are trademarks owned by the ProStrakan group of companies.

17.5 FDA-Approved Patient Labeling

Patient Information

Sancuso® [san-KOO-so]
(Granisetron Transdermal System)

IMPORTANT: For skin use only


Read the Patient Information that comes with Sancuso before you start using it and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment. If you have any questions about Sancuso, ask your healthcare provider.

What is Sancuso?

Sancuso is a prescription medicine used to prevent nausea and vomiting in people receiving some types of chemotherapy treatment.

Sancuso is a skin patch that slowly releases the medicine contained in the adhesive (glue), through clean and intact skin areas into your bloodstream while you wear the patch.

Important: Sancuso contains granisetron, the same medicine in Kytril. Do not take Kytril at the same time you use Sancuso unless your healthcare provider tells you it is alright.



Who should not use Sancuso?

Do not use Sancuso if you are allergic to any of the ingredients in Sancuso. See the end of this leaflet for a list of ingredients in Sancuso.

What should I tell my healthcare provider before using Sancuso?

Tell your healthcare provider about all your medical conditions, including if you:

- are allergic to medical adhesive tape, adhesive dressings or other skin patches.
- have pain or swelling in your stomach area (abdomen).
- are pregnant. It is not known if Sancuso will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant.
- are breast-feeding or plan to breast-feed. It is not known if Sancuso passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and non-prescription medicines, vitamins and herbal supplements. Other medicines may affect how Sancuso works. Sancuso may also affect how other medicines work.

How should Sancuso be used?

Use Sancuso exactly as prescribed.

See the detailed Patient Instructions for Applying Sancuso at the end of this Patient Information leaflet.

What should I avoid while using Sancuso?

Avoid sunlight. The medicine in Sancuso (granisetron) may not work as well and/or may affect your skin if exposed to direct sunlight or the light from sunlamps or tanning beds. It is important to do the following:

- While you wear the patch, keep it covered with clothing if you will be in sunlight or near a sunlamp, including tanning beds.
- Keep the skin where Sancuso was applied covered for another 10 days after the patch is taken off to protect from exposure to direct sunlight.

What are the possible side effects of Sancuso?

Sancuso can cause serious side effects:

- Using Sancuso may make it harder to identify certain stomach and bowel problems that are from other causes.** Tell your healthcare provider if you have any stomach area (abdominal) pain or swelling while using Sancuso.
- Skin reactions.** Skin reactions can happen just at the patch application site or outside the patch application site. Tell your healthcare provider if you get any redness, rashes, bumps, blisters or itching at the patch application site, and especially if they spread outside the place where the patch was or if they appear outside the patch application site. You may need to stop using Sancuso.

Common side effects of Sancuso are:

- constipation
- headache.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Sancuso. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Sancuso?

- Keep Sancuso in the package it comes in.
 - Store Sancuso at 20-25°C (68-77°F).
- Keep Sancuso out of the reach of children.**

General information about Sancuso

Medicines are sometimes prescribed for conditions that are not mentioned in Patient Information leaflets. Do not use Sancuso for a condition for which it is not prescribed. Do not give Sancuso to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information leaflet summarizes the most important information about Sancuso. If you would like more information, talk to your healthcare provider. You can ask your pharmacist or healthcare provider for information about Sancuso that is written for health professionals.

For more information, go to www.sancuso.com or call 1-800-SANCUSO.

Patient Instructions for Applying Sancuso

When do I apply the Sancuso patch?

- Apply Sancuso at least 1 day (24 hours) before your scheduled chemotherapy treatment.
- You may apply Sancuso up to 2 days (48 hours) before your scheduled chemotherapy.
- Wear the patch all the time during your chemotherapy.
- Sancuso may be worn for up to 7 days, depending on how long your chemotherapy treatment lasts (up to 5 days).
- Remove the patch at least 1 day (24 hours) after your chemotherapy is finished.
- Keep the patch covered, such as under clothing, while you are wearing it to avoid a skin reaction to sunlight or sunlamps. Keep the skin where Sancuso was applied (application site) covered up for another 10 days after the patch is taken off to prevent a skin reaction. See "What should I avoid while using Sancuso?"

Where do I apply the Sancuso patch?

- Apply Sancuso to a clean, dry, healthy area of skin on the outside part of your upper arm.



- The area you choose should not be oily, recently shaved or have any skin problems such as being damaged (cut or scraped) or irritated (redness or a rash).
- Do not apply Sancuso to areas that have been treated with creams, oils, lotions, powders or other skin products that could keep the patch from sticking well to your skin.**


How do I apply the Sancuso patch?

The Sancuso patch comes inside a pouch which is inside the carton.

- Do not remove the patch from the pouch until you are ready to use it.
- Do not cut the Sancuso patch into pieces.**
- Remove the pouch from the carton.




- Tear the pouch open using the slit provided, and remove the patch. Each pouch contains one Sancuso patch stuck onto a rigid plastic film, and a separate thin, clear protective liner.



- Remove the thin, clear protective liner to expose the printed side of the patch. Throw away the liner. The protective liner is only included in the pouch to separate the patch from the inside of the pouch, and is not part of the patch.



- The unprinted, sticky side of the patch is covered by a two-piece rigid plastic film. Bend the patch in the middle and remove one half of the rigid plastic film. Be careful not to stick the patch to itself and avoid touching the sticky side of the patch.



- While holding the remaining half of the rigid plastic film, apply the patch to your skin. Remove the second half of the rigid plastic film and press the whole patch firmly in place with your fingers and smooth down. Press firmly making sure it sticks well to the skin, especially around the edges.



- Wash your hands right away after applying the patch to remove any medicine that may have stuck to your fingers.
- Keep the patch in place for the whole time you are having chemotherapy. Remove the patch at least 1 day (24 hours) after your chemotherapy is finished. The patch can be worn for up to 7 days, depending on the number of days your chemotherapy treatment lasts.
- Do not re-use the patch after you remove it. See below for instructions on the right way to remove and throw away the patch.

What to do if the Sancuso patch does not stick well?

If the patch does not stick well, you may use surgical bandages or medical adhesive tape to keep the patch in place. Place tape or bandages on the edges of the patch. Do not completely cover the patch with bandages or tape and do not wrap completely around your arm. If the patch comes more than half off or it becomes damaged see your healthcare provider.

Can I bathe or shower while wearing Sancuso?

You can continue to shower and wash normally while wearing the Sancuso patch. It is not known how other activities, for example swimming, strenuous exercise or using a sauna or whirlpool, may affect Sancuso. Avoid these activities while wearing Sancuso.

How do I remove and dispose of Sancuso?

- When you remove the patch, peel it off gently.
- The used patch will still contain some of the medicine. After removing the used Sancuso patch, fold it in half so that the sticky side sticks to itself. Throw away the Sancuso patch in the garbage, out of the reach of children and pets. Do not re-use the patch.
- After removing the patch you may find some adhesive is left on your skin. Gently wash the area with soap and water to remove it. Do not use alcohol or other dissolving liquids, such as nail polish remover. These may cause skin irritation.
- Wash your hands after handling the patch.
- You may see mild redness on the skin where the patch is removed. This redness should go away within three days. If redness continues, tell your healthcare provider.

What are the ingredients in Sancuso?

Active ingredient: granisetron. Inactive ingredients: acrylate-vinylacetate copolymer, polyester, titanium dioxide, polyamide resin and polyethylene wax.