

Sancuso®
(Granisetron Transdermal System)

the
difference
between life
and living

1 patch 5 days
Prevention of CINV*

*CINV=Chemo-induced nausea and vomiting



Important Safety Information

What is SANCUSO (granisetron transdermal system)?

SANCUSO is a prescription medicine used to prevent nausea and vomiting in adults receiving anti-cancer (chemotherapy) treatment that causes moderate or severe vomiting.

SANCUSO is a skin patch (transdermal system) that slowly releases the medicine into your bloodstream while you wear the transdermal system.

It is not known if SANCUSO is safe and effective in children.

www.sancuso.com

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.



SANCUSO prevents nausea and vomiting caused by chemotherapy

Most patients expect to experience nausea and vomiting while undergoing chemotherapy. In fact, nausea and vomiting are among the greatest fears they have. Fortunately, with proper treatment, you do not have to suffer from these side effects.

SANCUSO (Granisetron Transdermal System) is the first and only patch approved by the US Food and Drug Administration (FDA) for the prevention of CINV in patients receiving some types of chemotherapy treatment. With nearly 2 decades of real-world use, SANCUSO is an easy-to-apply patch that continuously delivers medication to help prevent CINV for up to 5 full days.¹

This brochure will help you and your loved ones learn a little more about CINV and help you get ready for your treatment with SANCUSO.



Important Safety Information (continued)

Who should not use SANCUSO?

Do not use SANCUSO if you are allergic to granisetron or any of the ingredients in SANCUSO. See the end of this leaflet for a list of ingredients in SANCUSO.

What should I tell my healthcare provider before using SANCUSO?

Before using SANCUSO, tell your healthcare provider about all of your medical conditions, including if you:

- have pain or swelling in your stomach area (abdomen).
- are pregnant. It is not known if SANCUSO will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if SANCUSO passes into your breast milk.



www.sancuso.com

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.

1. Sancuso (package insert), Nashville, TN; Cumberland Pharmaceuticals Inc.

* CINV = Chemo-induced nausea and vomiting

What makes SANCUSO different?¹

SANCUSO is the only skin patch approved by the FDA for the prevention of CINV in patients receiving chemotherapy that is likely to cause nausea or vomiting.

There are no infusions needed or excessive pill regimens patients have to remember to follow.

The active drug in SANCUSO slowly dissolves in the thin layer of adhesive that sticks to your skin and is released into your bloodstream, working continuously to prevent CINV.

SANCUSO is applied 24 to 48 hours before receiving the first dose of chemotherapy and can prevent CINV for up to 5 consecutive days (it may be worn for up to 7 days total).



Important Safety Information (*continued*)

What should I tell my healthcare provider before using SANCUSO?

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. Other medicines may affect how SANCUSO works. SANCUSO may also affect how other medicines work. SANCUSO contains granisetron. Do not take other granisetron containing products with SANCUSO.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I use SANCUSO?

- Read the Instructions for Use that comes with SANCUSO transdermal system.
- Use SANCUSO exactly as your healthcare provider tells you to.

www.sancuso.com

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.

How SANCUSO helps prevent CINV



From the time that you apply the patch, SANCUSO slowly and continuously delivers granisetron through your skin and into your bloodstream until the patch is removed within 7 days.



Granisetron blocks cells in the small intestine from sending messages to the brain that trigger CINV.



Granisetron blocks cells in one area of the brain from sending messages to a different area of the brain that triggers CINV.

Because the brain does not receive these messages, feelings of nausea and/or vomiting are less likely to occur.

1. Sancuso (package insert), Nashville, TN; Cumberland Pharmaceuticals Inc.

* CINV = Chemo-induced nausea and vomiting

Wearing SANCUSO



Avoid sunlight. The medicine in SANCUSO may not work as well and/or may affect your skin if exposed to direct sunlight or the light from sunlamps or tanning beds. If there is a risk of exposing the patch to sunlight or sunlamps, you must keep it covered under clothing. Continue to keep the patch area covered for 10 days after removing SANCUSO.



Avoid heat sources. Avoid exposing SANCUSO and the surrounding area to direct external heat sources, such as heating pads.



Showering/bathing. You can continue to shower and bathe normally while wearing SANCUSO. However, you should try to avoid immersing the patch in water for long periods of time to keep it from coming off.



Avoid certain activities. It is not known whether activities such as swimming, strenuous exercise, or use of a sauna or whirlpool may affect the patch. Therefore, you should also avoid these activities while wearing SANCUSO.



Important Safety Information (continued)

What should I avoid while using SANCUSO?

Do not apply any heat source over or near the SANCUSO transdermal system.

- A heating pad or heat lamp should not be used where the transdermal system is applied.
- You should avoid long periods of exposure to heat because heat can increase the amount of SANCUSO in your blood.

www.sancuso.com

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.

Possible side effects of SANCUSO

SANCUSO can cause side effects in some patients. The most common side effect with SANCUSO is constipation.

Be sure to notify your healthcare professional if you have pain in your abdomen or your abdomen becomes swollen.

While you are wearing SANCUSO, you may see some mild skin redness at or around the patch. Call your healthcare professional if you see any redness, especially if it spreads outside the area of skin where SANCUSO is applied.

When not to use SANCUSO

Avoid use if you are allergic to granisetron or any of the other ingredients in SANCUSO, or if you have allergies to medical adhesive tape, adhesive dressings, or other skin patches.

Avoid use if you are pregnant or breastfeeding unless your healthcare professional has told you that you can use it. Be sure to tell your healthcare professional if you are pregnant, if you become pregnant, or plan to become pregnant while using SANCUSO, or if you are planning to breastfeed or are breastfeeding.



Important Safety Information (continued)

What should I avoid while using SANCUSO?

Avoid sunlight and artificial sunlight. The medicine in SANCUSO may not work as well and may affect your skin if exposed to direct sunlight or artificial sunlight from sunlamps or tanning beds.

- Keep the transdermal system covered with clothing if you will be in direct sunlight or artificial sunlight.
- Keep the skin where SANCUSO was applied covered for another 10 days after the transdermal system is taken off to protect from exposure to direct sunlight or artificial sunlight.

* CINV = Chemo-induced nausea and vomiting



Important Safety Information (continued)

What are the possible side effects of SANCUSO?

SANCUSO may cause serious side effects, including:

- **Using SANCUSO may make it harder to identify certain stomach (abdomen) and bowel problems that are from other causes.** Tell your healthcare provider if you have any abdominal pain or swelling while using SANCUSO.
- **Serotonin Syndrome.** A potentially life-threatening problem called serotonin syndrome can happen if you use SANCUSO with certain medicines. Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the following signs and symptoms of serotonin syndrome:

• agitation	• sweating
• seeing or hearing things that are not real	• flushing (hallucinations)
• confusion	• high body temperature (hyperthermia)
• coma	• shaking (tremors), stiff muscles, or muscle twitching
• fast heart beat	• loss of coordination
• changes in blood pressure	• seizures
• dizziness	• nausea, vomiting, diarrhea
- **Skin reactions.** Skin reactions can happen at the transdermal system application site or outside the transdermal system application site. Tell your healthcare provider if you get any redness, rashes, bumps, blisters or itching at or near the transdermal system application site, and especially if they spread outside the place where the transdermal system was applied or if they appear outside the transdermal system application site. You may need to stop using SANCUSO.

Tips for applying

- Be sure to keep the SANCUSO patch inside the sealed pouch until you are ready to apply it
- Do not cut the SANCUSO patch into pieces
- Apply SANCUSO a minimum of 24 to a maximum of 48 hours before your scheduled chemotherapy treatment
- Apply SANCUSO to a clean, dry, nearly hairless area of skin on the outside of your upper arm. Do not put SANCUSO on areas that have been treated with creams, oils, lotions, powders, or other skincare products that might keep the patch from sticking to your skin. SANCUSO should not be placed on skin that is red, irritated, cut, or scraped
- If the patch does not stick well, you may use surgical bandages or medical adhesive tape to keep the patch in place. Place tape or bandages on the edges of the patch. Do not completely cover the patch with bandages or tape and do not wrap completely around your arm. If the patch comes more than half off or it becomes damaged, call your healthcare professional
- If you've attached your patch and your chemotherapy is delayed or canceled or if the patch completely falls off, the SANCUSO Patch Replacement Program may be able to help you get a replacement patch. Go to patientrxsolutions.com to learn more



www.sancuso.com


Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.

* CINV = Chemo-induced nausea and vomiting

Follow these steps for proper application


Apply to a clean, dry, healthy area of skin on the outside part of your upper arm

- Do not cut the SANCUSO patch into pieces




Remove the patch from the carton; tear the pouch open and remove the patch. Each pouch contains one SANCUSO patch stuck into a rigid plastic film, and a separate clear protective liner.

Do not remove the SANCUSO patch from the pouch until you are ready to use it



Remove the thin, clear protective liner to expose the printed side of the patch.



Bend the patch in the middle and remove one half of the rigid plastic film. Be careful not to stick the patch to itself and avoid touching the sticky side of the patch.



While holding the remaining half of the rigid plastic film, apply the patch to the skin. Remove the second half of the rigid plastic film and press the whole patch firmly in place with your fingers and smooth down. Press firmly, making sure it sticks well, especially around the edges.



Wash your hands right away after applying the patch to remove any medicine that might have stuck to your fingers.



Keep the patch in place for the duration of the chemotherapy. Remove the patch at least 1 day (24 hours) after chemotherapy is finished. The patch can be worn up to 7 days, depending on the number of days that your chemotherapy treatment lasts.

Do not reuse the SANCUSO patch after you remove it. See instructions included with your SANCUSO patch on how to remove and throw away after use



Important Safety Information *(continued)*

The most common side effect of SANCUSO is constipation.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of SANCUSO. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

www.sancuso.com

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.

* CINV = Chemo-induced nausea and vomiting

Eligible patients may save up to \$300 per patch on Sancuso after paying the first \$20 per prescription.^a



Here's How: Your healthcare professional prescribes the SANCUSO patch, a prescription medicine used to prevent CINV.



Your healthcare professional will submit your prescription information to Patient Rx Solutions, called the "HUB" team, at

www.patientrxsolutions.com



If you are eligible, the HUB team will work with your healthcare professional and your commercial insurance provider to help you receive the appropriate number of medically necessary SANCUSO patches that you are allowed to receive for a \$20 out-of-pocket cost.^a

Ask your healthcare professional today to register you for the Patient Rx Solutions program.

^aThe SANCUSO Patient Rx Solutions program is only for commercially insured patients. Please visit www.patientrxsolutions.com for complete terms and conditions, information about access to SANCUSO, or if you require other reimbursement assistance.

patient **Rx** solutions

Provides services and programs to make it easier for you to gain access to and pay for SANCUSO

Reimbursement Assistance Program. If you have prescription insurance coverage, our experienced staff will work directly with you for answers to all of your SANCUSO coverage-related questions.

Copay Assistance Card Program. The Sancuso Copay Assistance Card can be accessed at <https://sservices.trialcard.com/Coupon/Sancuso>. Eligible patients may save up to \$300 per patch on Sancuso after paying the first \$20 per prescription.^a

SANCUSO Patch Replacement Program. If your chemotherapy is delayed or canceled, the SANCUSO Patch Replacement Program may be able to help you get a replacement patch through your physician's office and sent directly to your home.

For additional information on any of these programs, please call 1-844-RXSOLUTIONS (1-844-797-6588) or visit www.patientrxsolutions.com.

* CINV = Chemo-induced nausea and vomiting

www.sancuso.com

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.



Important Safety Information

What is SANCUSO (granisetron transdermal system)?

SANCUSO is a prescription medicine used to prevent nausea and vomiting in adults receiving anti-cancer (chemotherapy) treatment that causes moderate or severe vomiting.

SANCUSO is a skin patch (transdermal system) that slowly releases the medicine into your bloodstream while you wear the transdermal system.

It is not known if SANCUSO is safe and effective in children.

Who should not use SANCUSO?

Do not use SANCUSO if you are allergic to granisetron or any of the ingredients in SANCUSO. See the end of this leaflet for a list of ingredients in SANCUSO.

What should I tell my healthcare provider before using SANCUSO?

Before using SANCUSO, tell your healthcare provider about all of your medical conditions, including if you:

- have pain or swelling in your stomach area (abdomen).
- are pregnant. It is not known if SANCUSO will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if SANCUSO passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. Other medicines may affect how SANCUSO works. SANCUSO may also affect how other medicines work. SANCUSO contains granisetron. Do not take other granisetron containing products with SANCUSO.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I use SANCUSO?

- Read the Instructions for Use that comes with SANCUSO transdermal system.
- Use SANCUSO exactly as your healthcare provider tells you to.

What should I avoid while using SANCUSO?

Do not apply any heat source over or near the SANCUSO transdermal system.

- A heating pad or heat lamp should not be used where the transdermal system is applied.
- You should avoid long periods of exposure to heat because heat can increase the amount of SANCUSO in your blood.

www.sancuso.com

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.

Avoid sunlight and artificial sunlight. The medicine in SANCUSO may not work as well and may affect your skin if exposed to direct sunlight or artificial sunlight from sunlamps or tanning beds.

- Keep the transdermal system covered with clothing if you will be in direct sunlight or artificial sunlight.
- Keep the skin where SANCUSO was applied covered for another 10 days after the transdermal system is taken off to protect from exposure to direct sunlight or artificial sunlight.

What are the possible side effects of SANCUSO?

SANCUSO may cause serious side effects, including:

- **Using SANCUSO may make it harder to identify certain stomach (abdomen) and bowel problems that are from other causes.** Tell your healthcare provider if you have any abdominal pain or swelling while using SANCUSO.

- **Serotonin Syndrome.** A potentially life-threatening problem called serotonin syndrome can happen if you use SANCUSO with certain medicines. Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the following signs and symptoms of serotonin syndrome:

• agitation	• sweating
• seeing or hearing things that are not real	• flushing (hallucinations)
• confusion	• high body temperature (hyperthermia)
• coma	• shaking (tremors), stiff muscles, or muscle twitching
• fast heart beat	• loss of coordination
• changes in blood pressure	• seizures
• dizziness	• nausea, vomiting, diarrhea

- **Skin reactions.** Skin reactions can happen at the transdermal system application site or outside the transdermal system application site. Tell your healthcare provider if you get any redness, rashes, bumps, blisters or itching at or near the transdermal system application site, and especially if they spread outside the place where the transdermal system was applied or if they appear outside the transdermal system application site. You may need to stop using SANCUSO.

The most common side effect of SANCUSO is constipation.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of SANCUSO. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Sancuso® the difference (Granisetron Transdermal System) **between life and living**

Prevent chemotherapy-induced nausea and vomiting for up to 5 full days¹

No pills to remember. No injections. Just one patch.
Ask your healthcare professional if SANCUSO is right for you.

Visit **www.sancuso.com** or call **1-833-SANCUSO** (1-833-726-2876).

Patient Rx Solutions provides services and programs to gain access to and pay for the medications you need^a:

- Reimbursement Assistance Program
- Copay Assistance Card Program
- SANCUSO Patch Replacement Program

patient **Rx**«
solutions

For additional information on any of these programs,
please call 1-844-RXSOLUTIONS (1-844-797-6588) or
visit www.patientrxsolutions.com.



Important Safety Information

What is SANCUSO (granisetron transdermal system)?

SANCUSO is a prescription medicine used to prevent nausea and vomiting in adults receiving anti-cancer (chemotherapy) treatment that causes moderate or severe vomiting.

SANCUSO is a skin patch (transdermal system) that slowly releases the medicine into your bloodstream while you wear the transdermal system.

It is not known if SANCUSO is safe and effective in children.

Please see additional Important Safety Information throughout and full Important Safety Information on pages 14-15. See accompanying Patient Information.

^a The various Patient RX Solutions programs are subject to eligibility criteria.

1. Sancuso (package insert), Nashville, TN; Cumberland Pharmaceuticals Inc. **www.sancuso.com**